



Influence of COVID-19 on Quality of Life among Nurses Treating COVID-19 in Tamil Nadu

Pooja Sadasivam^a, Preetha S^{a†}, G. Sridevi^{a‡} and Lavanya Prathap^{b‡}

^a Department of Physiology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai. India.

^b Department of Anatomy, Saveetha Dental college and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai. India.

Authors' contributions

This work was carried out in collaboration among all authors. Author PS Framing the manuscript, did Statics approval, approved the final draft of the manuscript. Author P. Sadasivam managed the literature collection. Authors GS and LP approved final version of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i58B34203

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78925>

Original Research Article

Received 10 October 2021
Accepted 14 December 2021
Published 15 December 2021

ABSTRACT

Background: COVID-19 is caused by the novel severe acute respiratory syndrome (SARS- COV-2). The coronavirus is transmitted by different ways, including contact transmission, direct transmission and aerosol transmission. In health care professions the largest and diverse forces are the nurses. Nurses are important in managing the health emergency management crisis because of their vital link between the health care professionals and the patients. The COVID-19 pandemic has placed health care professionals (HCPs) in high stressful circumstances due to increase in patients. Normal life of nurses would be unprecedented disruption and high risk of exposure. The aim of the study is to investigate the quality of life among nurses treating patients during the COVID-19 pandemic.

Materials and Methods: The study setting was an online questionnaire survey among nurses in Tamilnadu. This study was conducted in February 2021. There are 127 participants in this survey. The answers regarding how they feel treating COVID-19 Patients, kind of support, impact and quality of life were collected. The data were analysed statistically using SPSS software. The results and observations were recorded in a pie chart. The chi square test and correlation between genders was done and represented in the bar chart. $p < 0.05$ was statistically significant.

[†]Assistant Professor;

[‡]Associate Professor;

*Corresponding author: E-mail: preethas.sdc@saveetha.com;

Results: More than 50% of the nurses responded that their quality of life was good. 43.31% Females have a good quality of life. 78% of the nurses replied that they have both physical and mental impact treating COVID-19 Patients. The association between the gender and quality of life of nurses was found with the help of chi square test $p=0.383$ ($p >0.05$) and hence it was statistically not significant.

Conclusion: The present study found that the majority of nurses 54.33% have a good quality of life. The officials in the health care system should improve the nurses' working conditions, strategies and their quality of work life, so that nurses will perform their duties properly.

Keywords: Nurses; COVID-19; quality of life; health care professionals; SARS-COV-2; innovative techniques.

1. INTRODUCTION

COVID-19 is caused by the novel severe acute respiratory syndrome (SARS-COV-2). The coronavirus in humans will cause infection in the respiratory tract that ranges from mild to lethal [1]. The coronavirus is transmitted by different ways: contact transmission, direct transmission and aerosol transmission. In December 2019, an unknown case was admitted in Wuhan China with pneumonia [2]. In India on 30 January 2020, the first laboratory confirmed case of COVID -19 was reported. After that till now there is an exponential growth in cases. Many government and public health policies that were developed to decrease the spread of coronavirus and had adopted a population health approach [3].

Nurses played the most important role in preventing the infection, controlling the infection, Isolation, contamination and public health [4]. This COVID -19 pandemic caused stressful circumstances among the health care professionals (HCPs) due to the increase of patients' load [5]. In the 2003 SARS (severe acute respiratory syndrome) outbreak in China, one third of fatalities were health care professionals [6].

The front line workers feel a higher level of work intensity and get easily tired due to wearing protective equipment causing some inconvenience to breathing, drinking water and vision. Over 200 health care professionals including doctors and nurses have been reported in Tamilnadu with COVID-19. The WHO has stated that COVID-19 pandemic has caused high levels of anxiety, stress, and depression [7]. However, COVID-19 pandemic hits in India affected the mental health status of Indian health care professionals. The experiences from our previous studies [8] have led us to focus on the current topic. Our team has extensive knowledge and research experience that has translated into

high quality publications [9–28]. Therefore the aim of our study was to investigate the quality of life among nurses treating patients during the COVID-19 pandemic.

2. MATERIALS AND METHODS

The study setting was done through an online questionnaire survey among nurses in Tamilnadu. The advantage of online surveys is that it is time saving. There are 127 participants in this survey. This study was conducted in the mid month of February 2021. The lists of independent variables are age and gender and the dependent variable is quality of life among nurses. The survey included questions like age, gender, experience, satisfaction after treating, kind of support from others and friends, enough energy for everyday life, ability to perform daily life activities, fear of transmission, opportunity for leisure activities, feel safe, quality of life, safety measures and kind of impact they face. The survey was conducted through online Google forms and shared in a whatsapp groups among nurses. The response was transferred to MS Excel. The data were analyzed statistically using SPSS software. The results were recorded in the form of pie charts. The chi square test and correlation between gender was done and represented in the bar charts (P value less than 0.05 is statistically significant).

3. RESULTS

It was found that the majority of the nurses age between 20-30. In this 80.31% nurses are female and 19.69% nurses are male. Majority of nurses were married (63.41%). Among this 36.59% nurses' experience was 3years. 58.27% nurses satisfied with treating COVID-19 Patients (Fig. 1). Nurses of 49.61% have moderate levels of energy to do their everyday life activities. 54.33% of nurses replied that they have a good quality of life (Fig. 2). 43.90% nurses have a moderate

level of support from others and 55.26% nurses responded that they were satisfied with the kind of support from friends. 56.91% nurses were satisfied and they have the ability to perform daily life activities. 47.15% nurses have moderate levels of fear of transmission of disease to family members. 61.79% of nurses replied that they have little time for leisure activity. 43.05% felt that they were safe. Most of the nurses (74.80%) responded that they have enough safety measures. 77.95% nurses replied that they impacted both physically as well as mentally (Fig. 3). The association between the gender and quality of life of nurses was found with the help of chi square test $p=0.383$ ($p > 0.05$) and hence it was statistically not significant (Fig. 4). Similarly the association between the gender and ability to perform daily life activities was determined with the help of chi square test $p=0.511$ ($p > 0.05$) and hence it was statistically not significant (Fig. 5).

4. DISCUSSION

When the participants of the survey were questioned about the quality of life among nurses

treating COVID-19 patients, 54.33% of the participants responded that they have a good quality of life. Even during the COVID-19 pandemic period nurses felt good quality of life. Previous study conducted among the nurses before Covid 19 pandemic reported that 60% of the nurses had a moderate level of quality of working life while 37.1% and only 2% had good quality of life [29]. When the participants were questioned about whether they have any fear of transmission of COVID-19 to family members 47.15% of nurses have moderate levels of fear. In Previous study, 59.5% had fear that they might infect family members [30].

The participants responded to another question to what extent did nurses have opportunities for leisure activity, 61.79% of the nurses responded that they have only a little time for leisure activity. Previous study conducted among health care workers reported that during the early stage of the SARS epidemic health care workers have poor sleep and insomnia [31]. The present study also showed that the majority of nurses (74.80%) have enough safety measures to treat the COVID-19 patients. Even though health care

GRAPHS:




Fig. 1. This pie chart represents the work satisfaction of nurses treating COVID19 patients. Orange colour denotes dissatisfied, light green denotes neither satisfied nor dissatisfied, dark green denotes satisfied and Dark magenta denotes very dissatisfied. Most of the nurses (58.27%) responded that they were satisfied with treating COVID-19 Patients, 25.98% were dissatisfied, 13.39% were neither satisfied nor dissatisfied and 2.36% were very dissatisfied




Fig. 2. The pie chart represents the Quality of life among nurses. Spring green denotes good, red denotes poor, violet denotes very good and grey denotes neither poor nor good. Majority of nurses (54.33%) replied that they have a good quality of life, 26.77% have very good quality of life, 13.39% have poor quality of life and 5.51% have neither poor nor good quality of life




Fig. 3. The pie chart represents the kind of impact faced by nurses treating COVID-19 patients. Mari gold colour denotes both physical and mental, forest green denotes mental, light sky blue denotes the physical. Most of the nurses (77.96%) replied that they impacted both physically as well as mentally, 14.17% were impacted mentally and 7.87% were impacted physically

professionals working all over the world received a guideline to treat or handle the COVID -19 patients and an adequate supply of medicinal items for protection (including masks, suits and glasses) they worked with the highest level of pressure [32]. The limitations of this study are

that less number of participants and majority of nurses were female. The future scope of this study is involving more male nurses with increased no of population and the study was done in some other states also.




Fig. 4. This bar graph depicts the association between the gender and quality of life of nurses. X - axis represents gender and Y- axis, number of responses. Blue represents good, green represents poor, sandal colour represents neither poor nor good, violet colour represents very good and yellow represents very poor. Females (58) responded that they have a good quality of life. Chi square test, $p=0.383$ ($p > 0.05$) and it was statistically not significant




Fig. 5. The bar graph depicts the association between gender and ability to perform daily life activities of nurses. X - axis represents gender and Y- axis, number of responses. Blue represents dissatisfied, green represents neither satisfied nor dissatisfied, sandal colour represents satisfied and violet represents very dissatisfied. Females (60) were satisfied to perform daily life activity. Chi square test, $p=0.511$ ($p > 0.05$) and it was statistically not significant

5. CONCLUSION

The present study found that the majority of nurses 54.33% have a good quality of life. The wholehearted hard work and selfless service of the nurses rendered by the nurses is to be appreciated. The officials in the health care system should improve the nurses working conditions strategies and their quality of work life, so that nurses will be able to perform their duties properly.

CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

SOURCE OF FUNDING

The present study is funded by Dheepam electricals, Dharmapuri, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University.

ACKNOWLEDGEMENT

We thank all the participants and Saveetha Dental College for their support to conduct this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Princeton B, Santhakumar P, Prathap L. Awareness on Preventive Measures taken by Health Care Professionals Attending COVID-19 Patients among Dental Students [Internet]. Vol. 14, European Journal of Dentistry. 2020;S105–9. Available:<http://dx.doi.org/10.1055/s-0040-1721296>
2. Catton H. Global challenges in health and health care for nurses and midwives everywhere. *Int Nurs Rev.* 2020;67(1):4–6.
3. Sun S, Lin D, Operario D. Need for a population health approach to understand and address psychosocial consequences of COVID-19. *Psychol Trauma.* 2020;12(S1):S25–7.
4. Smith GD, Ng F, Li WHC. COVID-19: Emerging compassion, courage and resilience in the face of misinformation and adversity [Internet]. *Journal of Clinical Nursing.* 2020;29:1425–8. Available:<http://dx.doi.org/10.1111/jocn.15231>
5. Ramasubramanian S, Preetha S, Premavathy D, Prathap L. Awareness on Spread of Misinformation and its Effect on Public with Regard to COVID-19 [Internet]. *International Journal of Current Research and Review.* 2020;12:66–73. Available:<http://dx.doi.org/10.31782/ijcrr.2020.sp08>
6. Hung LS. The SARS epidemic in Hong Kong: What lessons have we learned? [Internet]. *JRSM.* 2003;96:374–8. Available:<http://dx.doi.org/10.1258/jrsm.96.8.374>
7. Kang L, Ma S, Chen M, Yang J, Wang Y, Li R, et al. Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: A cross-sectional study [Internet]., *Brain, Behavior, and Immunity.* 2020;87:11–7. Available:<http://dx.doi.org/10.1016/j.bbi.2020.03.028>
8. Shruthi M, Preetha S. Effect of Simple Tongue Exercises in Habitual Snorers [Internet]. *Research Journal of Pharmacy and Technology.* 2018;11:3614. Available:<http://dx.doi.org/10.5958/0974-360x.2018.00665.0>
9. Saraswathi I, Saikarthik J, Senthil Kumar K, MadhanSrinivasan K, Ardhanaari M, Gunapriya R. Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: A prospective longitudinal study. *PeerJ.* 2020;8:e10164.
10. Santhakumar P, Roy A, Mohanraj KG, Jayaraman S, Durairaj R. Ethanolic Extract of Capparis decidua Fruit Ameliorates Methotrexate-Induced Hepatotoxicity by Activating Nrf2/HO-1 and PPAR γ Mediated Pathways. *Ind J Pharm Educ.* 2021;55(1s):s265–74.
11. Nambi G, Kamal W, Es S, Joshi S, Trivedi P. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment

- of chronic non-specific low back pain: A randomized controlled study. *Eur J Phys Rehabil Med.* 2018;54(6):880–9.
12. Rajakumari R, Volova T, Oluwafemi OS, Rajesh Kumar S, Thomas S, Kalarikkal N. Grape seed extract-soluplus dispersion and its antioxidant activity. *Drug Dev Ind Pharm.* 2020;46(8):1219–29.
 13. Clarizia G, Bernardo P. *Diverse Applications of Organic-Inorganic Nanocomposites: Emerging Research and Opportunities: Emerging Research and Opportunities.* IGI Global. 2019;237.
 14. Prakash AKS, Devaraj E. Cytotoxic potentials of *S. cumini* methanolic seed kernel extract in human hepatoma HepG2 cells [Internet]. *Environmental Toxicology.* 2019;34:1313–9. Available: <http://dx.doi.org/10.1002/tox.22832>
 15. Tahmasebi S, Qasim MT, Krivenkova MV, Zekiy AO, Thangavelu L, Aravindhan S, et al. The effects of oxygen-ozone therapy on regulatory T-cell responses in multiple sclerosis patients. *Cell Biol Int.* 2021;45(7):1498–509.
 16. Wadhwa R, Paudel KR, Chin LH, Hon CM, Madheswaran T, Gupta G, et al. Anti-inflammatory and anticancer activities of Naringenin-loaded liquid crystalline nanoparticles in vitro. *J Food Biochem.* 2021 Jan;45(1):e13572.
 17. Vivekanandhan K, Shanmugam P, Barabadi H, Arumugam V, Raj DDRD, Sivasubramanian M, et al. *Emerging Therapeutic Approaches to Combat COVID-19: Present Status and Future Perspectives* [Internet]. *Frontiers in Molecular Biosciences.* 2021;8. Available: <http://dx.doi.org/10.3389/fmolb.2021.604447>
 18. Ezhilarasan D. Critical role of estrogen in the progression of chronic liver diseases. *Hepatobiliary Pancreat Dis Int.* 2020;19(5):429–34.
 19. Egbuna C, Mishra AP, Goyal MR. *Preparation of Phytopharmaceuticals for the Management of Disorders: The Development of Nutraceuticals and Traditional Medicine.* Academic Press. 2020;574.
 20. Kamath SM, Manjunath Kamath S, Jaison D, Rao SK, Sridhar K, Kasthuri N, et al. In vitro augmentation of chondrogenesis by Epigallocatechin gallate in primary Human chondrocytes - Sustained release model for cartilage regeneration [Internet]. *Journal of Drug Delivery Science and Technology.* 2020;60:101992. Available: <http://dx.doi.org/10.1016/j.jddst.2020.101992>
 21. Barabadi H, Mojab F, Vahidi H, Marashi B, Talank N, Hosseini O, et al. Green synthesis, characterization, antibacterial and biofilm inhibitory activity of silver nanoparticles compared to commercial silver nanoparticles [Internet]. *Inorganic Chemistry Communications.* 2021;129:108647. Available: <http://dx.doi.org/10.1016/j.inoche.2021.108647>
 22. Bharath B, Perinbam K, Devanesan S, AlSalhi MS, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT–29 colon cancer cells [Internet]. *Journal of Molecular Structure.* 2021;1235:130229. Available: <http://dx.doi.org/10.1016/j.molstruc.2021.130229>
 23. GowhariShabgah A, Ezzatifar F, Aravindhan S, Olegovna Zekiy A, Ahmadi M, Gheibihayat SM, et al. Shedding more light on the role of Midkine in hepatocellular carcinoma: New perspectives on diagnosis and therapy. *IUBMB Life.* 2021;73(4):659–69.
 24. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med.* 2019;48(4):299–306.
 25. R H, Hannah R, Ramani P, Ramanathan A, Jancy MR, Gheena S, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene [Internet]. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology.* 2020;130:306–12. Available: <http://dx.doi.org/10.1016/j.oooo.2020.06.021>
 26. J PC, Pradeep CJ, Marimuthu T, Krithika C, Devadoss P, Kumar SM. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study [Internet]. *Clinical Implant Dentistry and Related Research.* 2018;20:531–4. Available: <http://dx.doi.org/10.1111/cid.12609>
 27. Wahab PUA, Abdul Wahab PU, Madhulaxmi M, Senthilnathan P, Muthusekhar MR, Vohra Y, et al. Scalpel

- Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study [Internet]. *Journal of Oral and Maxillofacial Surgery*. 2018;76:1160–4. Available:<http://dx.doi.org/10.1016/j.joms.2017.12.020>
28. Mudigonda SK, Murugan S, Velavan K, Thulasiraman S, Krishna Kumar Raja VB. Non-suturing microvascular anastomosis in maxillofacial reconstruction- a comparative study. *Journal of Cranio-Maxillofacial Surgery*. 2020;48(6):599–606.
 29. Moradi T, Maghaminejad F, Azizi-Fini I. Quality of working life of nurses and its related factors [Internet]. *Nursing and Midwifery Studies*. 2014;3. Available:<http://dx.doi.org/10.17795/nmsjournal19450>
 30. Stojanov J, Malobabic M, Stanojevic G, Stevic M, Milosevic V, Stojanov A. Quality of sleep and health-related quality of life among health care professionals treating patients with coronavirus disease-19 [Internet]. *International Journal of Social Psychiatry*. 2020; 002076402094280. Available:<http://dx.doi.org/10.1177/0020764020942800>
 31. McAlonan GM, Lee AM, Cheung V, Cheung C, Tsang KWT, Sham PC, et al. Immediate and sustained psychological impact of an emerging infectious disease outbreak on health care workers [Internet]. *The Canadian Journal of Psychiatry*. 2007;52:241–7. Available:<http://dx.doi.org/10.1177/070674370705200406>
 32. Song X, Zhou Y, Rao W, Zhang X. Comparison of prevalence and risk factors of somatization between Chinese health care workers and non-health care workers during COVID-19 outbreak [Internet]. Available:<http://dx.doi.org/10.21203/rs.3.rs-269527/v1>

© 2021 Sadasivam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78925>*